

Middlesex University Research Repository

An open access repository of

Middlesex University research

<http://eprints.mdx.ac.uk>

Chan, Wai Kit, Wildeboer, Dirk ORCID logoORCID: <https://orcid.org/0000-0003-1298-7511>,
Garelick, Hemda ORCID logoORCID: <https://orcid.org/0000-0003-4568-2300> and Purchase,
Diane ORCID logoORCID: <https://orcid.org/0000-0001-8071-4385> (2018) Competition of As
and other Group 15 elements for surface binding sites of an extremophilic Acidomyces
acidophilus isolated from a historical tin mining site. Extremophiles, 22 (5) . pp. 795-809. ISSN
1431-0651 [Article] (doi:10.1007/s00792-018-1039-2)

Final accepted version (with author's formatting)

This version is available at: <https://eprints.mdx.ac.uk/24935/>

Copyright:

Middlesex University Research Repository makes the University's research available electronically.

Copyright and moral rights to this work are retained by the author and/or other copyright owners unless otherwise stated. The work is supplied on the understanding that any use for commercial gain is strictly forbidden. A copy may be downloaded for personal, non-commercial, research or study without prior permission and without charge.

Works, including theses and research projects, may not be reproduced in any format or medium, or extensive quotations taken from them, or their content changed in any way, without first obtaining permission in writing from the copyright holder(s). They may not be sold or exploited commercially in any format or medium without the prior written permission of the copyright holder(s).

Full bibliographic details must be given when referring to, or quoting from full items including the author's name, the title of the work, publication details where relevant (place, publisher, date), pagination, and for theses or dissertations the awarding institution, the degree type awarded, and the date of the award.

If you believe that any material held in the repository infringes copyright law, please contact the Repository Team at Middlesex University via the following email address:

eprints@mdx.ac.uk

The item will be removed from the repository while any claim is being investigated.

See also repository copyright: re-use policy: <http://eprints.mdx.ac.uk/policies.html#copy>

This is a post-peer-review, pre-copyedit version of an article published in Extremeophiles.
The final authenticated version is available online at: <https://doi.org/10.1007/s00792-018-1039-2>

Competition of As and other Group 15 elements for surface binding sites of an extremophilic *Acidomyces acidophilus* isolated from a historical tin mining site

Wai Kit Chan, Dirk Wildeboer, Hemda Garelick, Diane Purchase

Department of Natural Science, Faculty of Science and Technology, Middlesex University, The Burroughs, London NW4 4BT, UK.

Corresponding author email: d.purchase@mdx.ac.uk

Telephone: +44 (0)20 8411 5262

Acknowledgements

The authors wish to acknowledge the expertise and thank Professor Ajit Shah, Dr Leonardo Pantoja Munoz and Ms Manika Choudhury for advice in the MALDI-TOF MS analysis, technical support in the analytical and microbiological work, respectively. We also like to thank Greevor Tin Mine for providing access and permission to sample on the site.

Abstract

An arsenic-resistant fungal strain, designated WKC-1, was isolated from a waste roaster pile in a historical tin mine in Cornwall, UK and successfully identified to be *Acidomyces acidophilus* using matrix-assisted laser desorption/ionization time-of-flight/time-of-flight tandem mass spectrometry (MALDI-TOF/TOF MS) proteomic-based biotyping approach. WKC-1 showed considerable resistance to As^{5+} and Sb^{5+} where the minimal inhibitory concentration (MIC) were 22500 mg L^{-1} and 100 mg L^{-1} respectively on Czapex-Dox Agar (CDA) medium; it was substantially more resistant to As^{5+} than the reference strains CBS 335.97 and CCF 4251. In a modified CDA medium containing 0.02 mg L^{-1} phosphate, WKC-1 was able to remove 70.30 % of As^{5+} (100 mg L^{-1}). Sorption experiment showed that the maximum capacity of As^{5+} uptake was 170.82 mg g^{-1} dry biomass as predicted by the Langmuir model. The presence of Sb^{5+} reduced the As^{5+} uptake by nearly 40%. Based on the Fourier-transform infrared spectroscopy

(FT-IR) analysis, we propose that Sb is competing with As for these sorption sites: OH, NH, CH, SO₃ and PO₄ on the fungal cell surface. To our knowledge, this is the first report on the impact of other Group 15 elements on the biosorption of As⁵⁺ in *Acidomyces acidophilus*.

Keywords

Acidomyces acidophilus, arsenic pollution, biosorption, bioremediation, MALDI-TOF/TOF-MS

Introduction

Due to the legacy of coal, tin and precious metals mining, abandoned mines constitute one of the most significant pollution hazards in Great Britain (Hudson-Edwards *et al.*, 2008). Mining operations disposed residues, often with high levels of transitional metals and metalloids, in the mining sites and these were often dispersed by water and/or wind resulting in far-reaching pollution concerns (Asklund and Eldvall, 2005; Wang and Mulligan, 2006). The major sources for transitional metals and metalloids in the mining industries are milling, grinding, concentrating ores, disposal of tailings operations as well as milling wastewater discharge (Adriano, 1986; Razo *et al.*, 2004). Roaster piles, tailing ponds and waste rock piles were some of the wastes left behind after mining operations ceased. Constant piling of such mine wastes resulted in an elevation of transitional metals and metalloids concentrations in the surrounding areas. The high soil contents of arsenic (As), iron (Fe), antimony (Sb) and zinc (Zn) and these have significant effect on the flora and fauna as well as human health (Dos Santos *et al.*, 2013).

In natural environments, compounds of metalloids such as As and Sb are widely dispersed as a consequence of anthropogenic and geological activities. As and Sb are by-products of tin-mining activities during the smelting process, where As is primarily found in the arsenopyrite (FeAsS) form (Telford *et al.*, 2009). The continuous disposal of arsenic trioxide (As₂O₃), a by-product in the furnace channel during the roasting process in tin mining activities, has been reported to cause serious contamination to surrounding soils and waters in proximity of mining sites. For instance, the concentration of As in soil adjacent to the Ron Phibun district tin mine in the Nakorn Si Thammarat province of southern Thailand was reported to be as high as 11000 mg kg⁻¹ (Francesconi *et al.*, 2002). To fully appreciate the toxic effects transitional metals exert on biological systems, it is important to analyse their bioavailability by determining the uptake of these metals from soil by microorganisms within a given time span (Olaniran *et al.*, 2013).

Geevor Tin Mine is a disused historical mine in the St Just mining district, one of the oldest mining districts in Cornwall (Yim, 1981). This tin mine was first established in the 1910s but due to the low global demand for tin and the high cost of operations, it was closed down in early 1990s. Upon closure of the mine, all the waste piles were abandoned on the site, the soil pollutants were contained and access to the site was restricted. According to Pirrie *et al.*, (2002), transitional metal and metalloids contamination is very common in Cornwall and it was estimated that approximately 1000 km² of Southwest of England are still contaminated with elevated concentrations of toxic metals and arsenic (Abrahams and Thornton, 1987; Camm *et al.*, 2004; Van Veen *et al.*, 2016). Metals bioavailability analysis of these soil samples will help to fully understand the actual amount of these metals available for uptake by microorganisms and their toxicity (Olaniran *et al.*, 2013).

Biosorption using fungal biomass has been receiving attention from many researchers globally as an alternative method to remove heavy metal/metalloid(s) from contaminated water and soil. It offers many advantages such as high efficiency, reduced operating cost, minimal usage of chemicals and low production of toxic chemical sludge (Gadd, 2009; Vijayaraghavan *et al.*, 2006). The transitional metals and metalloids present in the soil can be either already available or made available for uptake by microorganisms or plants, where they will be accessible for the sorption process (Peijnenburg and Jager, 2003; Del Giudice *et al.*, 2013; Antonucci *et al.*, 2017). It has also been established that ions from the same group in the periodic table could compete with each other during the biosorption process (Tsezos *et al.*, 1996).

A number of extremophile fungi have been successfully isolated from adverse environmental conditions. One of the most well-known is *Acidomyces acidophilus*, formerly known as *Scytalidium acidophilum*, and also known as the black fungi. It is a pigmented ascomycete capable of growing in extremely acidic conditions (Sigler and Carmichael, 1974). Its melanin-containing cell walls offer the fungus protection from adverse environmental conditions such as extreme pH, temperature and toxins (Jacobson *et al.*, 1995; Martin *et al.*, 1990; Tetsch *et al.*, 2006; Hujšlová *et al.*, 2013). This protection also provides the fungus a certain level of resistance to oxidative stress (Jung *et al.*, 2006). The enzymes produced by this fungus are of great interests as they can function at low pH and high temperatures and could have potential applications for a variety of industries (Polizeli *et al.*, 2005; Hess, 2008; Selbmann *et al.*, 2008). So far, there are no reports on the use of *A. acidophilus* for metalloids bioremediation.

This paper reports the isolation and characterisation of a highly resistant *A. acidophilus* WKC-1 strain from the disused mine in Cornwall that can tolerate high levels of As^{5+} . The ability of this isolate to remove As^{5+} is being investigated and sorption analyses carried out to determine its maximum adsorption capacity. The influence of Sb^{5+} and PO_4^{3-} on this isolate's capacity to remove As^{5+} has been studied to provide a better understanding of the relationship between its As-resistance and the presence of other chemicals in soil. Finally, the potential of using resistant fungi to bioremediate metalloids from polluted soil in historical sites is discussed.

Materials and Methods

Site description

The Geevor Tin Mine is located in the St Just District, Cornwall at 50°09' 06.43" N 5°40' 34.96" S, in the Southwest of England. It was the only tin mining site in the district after the closure of Levant Mine in 1930 (Noall, 1973) and ceased its operation in 1991 (Camm *et al.*, 2003). The site covers an area of 67 acres (270,000 m²) and it is now on the European Route of Industrial Heritage sites, an important tourist attraction in Cornwall.

Soil sampling

Six sampling points were selected as shown in Figure S1. Approximately 1 kg of surface soil samples from a depth of up to 0.5 m were collected randomly from each sampling point into sampling bags using a sterile trowel and spade. The soil samples were transported in an insulated cool box at 4 °C back to the laboratory within 24 h and stored in a refrigerator.

Soil analysis

Soil samples were air-dried for 72 h, ground finely using a pestle and mortar and sieved through a 2 mm sterile mesh prior to analysis. The pH of the soil samples, suspended in deionised water (soil:deionised water 1:2 w/v), was measured using a calibrated pH meter (Jenway, Model 3505). The soil organic matter (OM) content was determined using the ASTM (American Society of Testing and Materials) standard procedure (ASTM, 2000) and the cation exchange capacity (CEC) was analysed using the protocol recommended by Gillman and Sumpter (1986). The concentrations of As and Sb in each of the six soil samples were analysed using a three-step sequential extraction method for exchangeable (F1), weakly bound organic bound (F2) and residual (F3) fractions (Carapeto and Purchase, 2000). All the extracts (F1, F2 and F3) were

analysed using inductively coupled plasma optical emission spectrometry iCAP 1600 (ICP-OES) and the operating parameters summarised in Table S1. All the analyses were carried out in triplicates and the ICP-OES generated three readings per analysis. The percentage of bioavailability of both As and Sb was calculated by division of the summed fractions 1 and 2 by the total (F1+F2+F3) of each metalloid from the three-step sequential extraction. For analytical accuracy, the percentage recoveries (R) of all soil trace elements of interest were performed in soil certified reference materials (CRM) (#SQC001, lot 011233 and lot 017309, RTC, Laramie, WY, USA).

Enumeration and isolation of arsenic-tolerant fungi

Soil samples containing high As and Sb concentrations were used for screening of arsenic-resistant fungi. A ten-fold serial dilution was carried out using one gram of soil sample and plated out on to 2% malt extract agar (MEA; Oxoid Ltd., UK), supplemented with 100 mg L⁻¹ of chloramphenicol to prevent bacteria growth. The inoculated plates were incubated for 7 – 21 days at 25 °C and fungal viable counts determined. Colonies were sub-cultured, purified by passaging for ten times, screened in 2 % MEA at pH 1 containing As⁵⁺ (1000 – 25000 mg L⁻¹), prepared from sodium arsenate (Na₂HAsO₄). Fungal strains that survived the highest As-stress were considered as a potential candidate and maintained using the same MEA conditions with or without 100mg L⁻¹ of As⁵⁺.

Molecular identification of isolated fungi

Fungal isolates were grown on Potato Dextrose Agar (PDA) (CM0139, Oxoid Ltd, UK), pH 1 at 25 °C for 21 days. Mycelia were collected by pipetting Triton X-100 on the same colony spot for several times and transferring into a sterile tube. DNA was extracted using cetyl trimethylammonium bromide (CTAB) following the protocol by Stirling (2003) with a minor modification, DNA extraction was carried out twice on the samples at 65 °C for 50 min followed by bead milling. The extracted DNA was dissolved in 20 µl ultrapure water and stored at 4 °C.

The internal transcribed spacer (ITS) nuclear region of 18S-ITS1-5.8S-ITS2-28S rRNA of the fungal isolate was amplified by PCR using three sets of primers based on published sequences (White *et al.*, 1990; Martin and Rygiewicz, 2005). The first PCR used ITS1 forward and ITS2 reverse primers, the second used ITS5 forward and ITS4 reverse primers and the third used ITS1 forward and ITS4 reverse primers, all were obtained from Sigma-Aldrich and PCR amplifications performed (ITS1-ITS2 PCR: 94 °C for 2 min; 30 cycles of 94 °C for 1 min, 63 °C for 2 min, 72 °C for 1 min; followed by 72 °C for 10 min; ITS5-ITS4 and ITS1-ITS4 PCRs: 94 °C for 4 min;

159 30 cycles of 94 °C for 1 min, 58 °C for 1 min, 72 °C for 1 min; followed by 72 °C for 10 min).
 160 The PCR products were analysed by 2.0% (w/v) agarose gel electrophoresis and capillary
 161 electrophoresis on the MCE-202 MultiNA system (Shimadzu) in “on-tip analysis” mode
 162 following the protocol recommended by the manufacturer using the DNA 1000 reagent kit
 163 (Shimadzu) to quantify their concentrations and to confirm the results of the agarose gel
 164 electrophoresis. The PCR products were sequenced by GATC biotech (London, UK) and
 165 sequences analysed by nucleotide BLAST (NCBI) analysis. Based on the generated DNA
 166 sequences of the isolate and other reference sequences of fungi obtained from NCBI databases,
 167 a phylogenetic dendrogram from the evolutionary distance via the neighbour-joining method was
 168 constructed using bootstrap method of 1000 replications using Molecular Evolutionary Genetics
 169 Analysis 6 (MEGA6) software (Tamura *et al.*, 2013).

170 **Proteomics identification of fungal strain using MALDI-TOF/TOF MS**

171 Three reference strains of *A. acidophilus* were obtained from Centraalbureau Schimmelcultures
 172 (CBS), Netherlands (strain CBS 335.97) and Culture Collection of Fungi (CCF), Czech Republic
 173 (strains CCF4251 and CCF3679). Reference strains and the isolated fungal strain were grown in
 174 liquid salt medium (LSM) containing 2% dextrose, 0.1% (NH₄)₂SO₄, 0.001% K₂HPO₄, 0.05%
 175 MgSO₄·7 H₂O, 0.0026% FeSO₄ and 0.008% CaCl₂ with a final pH of 4.0-4.2 using a horizontal
 176 rotator (SB2 rotator, Stuart) for 72 h at room temperature. The sample preparation and extraction
 177 of proteins and peptides of *A. acidophilus* and the three reference strains were performed
 178 according to the Bruker fungi sample preparation protocol. Each extracted sample was analysed
 179 using a MALDI ground steel plate and six different sample spots (replicates) to generate six
 180 combined mass spectra (MSP) per fungal isolate. The reference strains for *A. acidophilus* CBS
 181 335.97, CCF4251 and CCF3679 were analysed to generate reference spectra and used to create
 182 an in house supplementary new database library for *A. acidophilus* fungal strains identification.
 183 The identification of the isolated fungal strain through comparison with reference strains and
 184 visualization of the mass spectra was performed with MALDI Biotyper software 3.0 (Bruker
 185 Daltonics).

186 **Determination of As minimum inhibitory concentration (MIC)**

187 The MIC for the isolated fungal strain and two *A. acidophilus* reference strains (CBS 335.97 and
 188 CCF 4251) were determined using solid acidic culture medium of modified Czapek dox agar
 189 (CDA) with either 1 mg L⁻¹ or 100 mg L⁻¹ PO₄³⁻ at pH 1, containing As⁵⁺ concentrations ranged
 190 from 1000 to 25000 mg L⁻¹. To allow polymerization of agar in culture medium at pH 1, double
 191 concentration of agar was added, and pH was adjusted after sterilization. The fungal mycelia

plugs were removed using a sterile pipette and placed in the middle of the agar. The plates were incubated for 21 days at 25 °C and the diameter of the each of the fungal colony was measured and MIC calculated from the average of the triplicate results.

Analysis of pH-effect on *A. acidophilus* WKC-1 growth

The effect of pH on the isolated *A. acidophilus* WKC-1 was determined using solid culture medium of MEA containing 1000 mg L⁻¹ of As⁵⁺ with pH ranged from 0.5 to 5, the desired pH was adjusted using NaOH (0.1M) or HCl (0.1M). The plates were incubated for 21 days at 25 °C and the diameter of the each of the fungal colony was recorded.

Arsenic removal efficiency

The efficiency of arsenic removal by *A. acidophilus* WKC-1 and three *A. acidophilus* reference strains were studied in a 0.15 mL centrifugal tube using LSM containing 1 g L⁻¹ of viable wet fungal biomass in pH 1 and supplemented with 100 mg L⁻¹ As⁵⁺, the cultures were cultivated using a horizontal rotator (SB2 rotator, Stuart) at 120 rpm for 21 days at room temperature and the final concentration As⁵⁺ in each filtrate was measured every 7 days using ICP-OES. All experiments were carried out in triplicates. The arsenic removal efficiency by all studied *A. acidophilus* strains was calculated using the following equation:

$$R = [(C_i - C_f) / C_i] \times 100$$

where:

R = Percentage As⁵⁺ removal;

C_i = Initial concentration of As⁵⁺ (mg L⁻¹);

C_f = Final concentration of As⁵⁺ (mg L⁻¹) after 21 days.

Biosorbent preparation and analysis of As biosorption

The *A. acidophilus* WKC-1 was inoculated in LSM for 21 days at 25 °C with constant shaking at 110 rpm using an orbital shaker (Minitron, Infors HT). The fungal biomass was harvested by filtration through Whatman No.11 filter paper, cleaned three times with deionised water to ensure the removal of all the excessive media residuals, freeze-dried (ScanVac CoolSafe, Labogene) for 24 h and grounded in mortar and pestle to fine powder. Each of the 1000 mg L⁻¹ As⁵⁺ and Sb⁵⁺ stock solution was prepared by dissolving Na₂HAsO₄ × 7 H₂O and KSb(OH)₆ (Sigma-Aldrich) in deionised water.

All adsorption tests were carried out in 50 mL conical flasks containing 20 mL of As⁵⁺ and/or Sb⁵⁺ solution at 25 °C on an orbital shaker at 120 rpm. Biosorption isotherms were formulated

through investigating the effect of pH and biomass loading capacity on the fungal cell as previously performed by Xu *et al.* (2012). In order to identify the pH effect on As⁵⁺ biosorption, two sets of experiments were carried out. Firstly, biosorption using fixed 0.5 g L⁻¹ dried fungal biomass in a range of As⁵⁺ (100 - 600 mg L⁻¹) and different pH range (1.0 - 6.0) was examined. Secondly, biosorption of a range of fungal biomass (0.5 - 5 g L⁻¹) using fix concentration of As⁵⁺ (500 mg L⁻¹) at different pH range was investigated. In order to investigate optimum contact time for the biosorption of As⁵⁺, samples were collected at different times (5, 15, 30, 60, 120 and 180 min) and filtered through Whatman No.11 filter paper. All filtrates were analysed for residual of As⁵⁺ concentration using ICP-OES. The uptake of As⁵⁺ by *A. acidophilus* WKC-1 was calculated using the following equation:

$$q_{eq} = \frac{V (c_i - c_{eq})}{m}$$

where:

- q_{eq} = As⁵⁺ uptake in mg per g biomass;
- V = Volume of As⁵⁺ used in mL;
- c_i = Initial concentration of As⁵⁺ (mg L⁻¹);
- c_{eq} = Equilibrium concentration of As⁵⁺ (mg L⁻¹);
- m = Amount of dry biosorbent (g).

FT-IR studies

The detection of vibration frequency changes in *A. acidophilus* WKC-1 for the untreated and As/Sb-treated biomass samples before and after the As⁵⁺ and Sb⁵⁺ biosorption were analysed using Fourier transform infrared spectroscopy (Travel IR, Perkin Elmer) and the attenuated total reflection (ATR) technique in the same experimental conditions as described for the biosorption experiment. Each biomass was freeze-dried and was placed on the single reflection diamond ATR crystal. The infra-red spectra were collected using the ATR-FTIR ranged from 400 to 4000 cm⁻¹ (Guibaud *et al.*, 2003).

Statistical analyses

All the experiments were performed in triplicates and the data obtained were calculated as mean plus/minus standard errors (mean ± SE). The statistical analysis on the difference in percentage of the bioavailability of As and Sb in soil samples was performed using 2-sample t-test. The As removal was compared between the isolated *A. acidophilus* WKC-1 strain and two *A. acidophilus* reference strains (CBS 335.97 and CCF 4521) using analysis of variance (one-way ANOVA). All the statistical analyses were performed using Minitab version 16.

Results

Physical and chemical properties of the soil samples

Results showed that all of the soil samples were acidic and the pH range varied sites with soil sample from site 3 being the most acidic (pH 1.13) and site 5 being the least acidic (pH 5.25). The highest and lowest CEC were 32.14 ± 4.31 meq/100g for soil sample 3 and 12.09 ± 3.71 meq/100g for soil sample 4 respectively, the mean CEC for all soil samples was 19.96 ± 3.57 . The highest and lowest OM contents were observed in soil samples 1 with 14.60 % and 4 with 3.34 % respectively. Results for the chemical and physical characterisation of the soil samples are summarised in Table 1.

Quality control data on the recovery of metal/metalloid(s) is shown in Table S2. The results showed good recovery with percentage recovery of As and Sb of more than 92% and 84% respectively using the two different certified metals in soil reference materials.

Table 1: Chemical and physical characteristics of soil samples from Geevor Tin Mine.

Site	Textural class	pH	% OM	CEC (meq/100 g)
1	Fine sand	3.75 ± 0.14	14.60%	19.85 ± 2.79
2	Fine sand	3.11 ± 0.07	13.55%	25.72 ± 4.16
3	Medium sand	1.13 ± 0.06	7.15%	32.14 ± 4.31
4	Clay	5.22 ± 0.12	3.34%	12.09 ± 3.71
5	Coarse sand	5.25 ± 0.09	10.68%	13.74 ± 3.08
6	Medium sand	3.26 ± 0.12	4.40%	16.22 ± 3.44

The total metal/metalloid concentration analysis showed that As levels exceeded those of Sb in all sampling sites (Table 2). The highest concentrations of As ($18043.50 \text{ mg kg}^{-1}$) and Sb ($213.69 \text{ mg kg}^{-1}$) were detected in soil sample 3, collected from the location of the roaster pile. Arsenic levels in all soil samples exceeded the UK Category 4 Screening Levels (C4SL) for commercial site of 640 mg kg^{-1} (Defra, 2014). The As concentration in soil sample from site 3 (obtained from the main roaster dump pile) exceeded the C4SL for the organically bound fractions (4511 mg kg^{-1}).

¹; Table 2). Since currently there is no C4SL values for Sb, the Dutch Guideline intervention values for soil remediation (Dutch Environment Ministry, 2013) was used to assess the extent of contamination (15 mg kg^{-1}). Only two soil samples (site 2 and site 3) were found to exceed the intervention limit.

The average percentages bioavailability of As and Sb in the soil samples from all six sites are presented in Figure 1. The bioavailability of As and Sb in all the soil samples were below 50%. However, the percentage of bioavailability of As was significantly higher than Sb in soil sample collected at site 3.




Figure 1: Percentage bioavailability of arsenic and antimony of soil from the sampling sites obtained from the summation of fraction 1 and 2 of the three-step sequential extraction.

Table 2: Mean As and Sb concentration ($\text{mg kg}^{-1} \pm \text{standard error}$) in the soil samples from the sampling sites using the three-step sequential extraction method. Data shown are the mean of three replicates.

Element					As				Sb			
Site	F1	F2	F3	Σ(F1 + F2 +F3)	F1	F2	F3	Σ(F1 + F2 +F3)				
1	1.13 ± 0.17	300.48 ± 7.41	1872.39 ± 44.22	2174.00 * +	0.10 ± 0.02	2.34 ± 0.88	4.65 ± 1.01	7.09				
2	0.66 ± 0.09	485.20 ± 1.54	5157.31 ± 17.38	5643.17 * +	0.96 ± 0.09	14.24 ± 2.19	32.4 ± 2.08	47.60 +				
3	249.20 ± 6.00	4511.30 ± 78.70	13283.00 ± 42.80	18043.50 * +	1.33 ± 0.03	13.96 ± 1.03	198.40± 4.03	213.69 +				
4	2.17 ± 0.08	325.60 ± 16.79	822.88 ± 21.02	1150.65 * +	0.12 ± 0.02	0.923 ± 0.14	3.067 ± 0.17	4.11				
5	11.64 ± 0.13	298.24 ± 5.10	1488.13 ± 18.98	1598.04 * +	ND	1.01 ± 0.22	2.35 ± 0.087	3.36				
6	0.28 ± 0.07	200.98 ± 3.81	955.82 ± 5.41	1158.88 * +	ND	2.39 ± 0.64	8.43 ± 0.66	10.82				

F1 Fraction 1 (exchangeable fraction), *F2* Fraction 2 (organically bound fraction), *F3* Fraction 3 (residual fraction), *ND* not detectable.

* Indicates exceeded the guideline values set by UK C4SL (for commercial site) and $^+$ indicates exceeded the intervention value limit set by the Dutch Guideline (Dutch Environment Ministry., 2013).

285 Identification and isolation of fungal strains

286 A total of 31 strains were isolated from soil samples collected from six different locations were
 287 exposed to As^{5+} ranged from 1000 to 22500 mg L^{-1} . Only one fungus from the most acidic and
 288 polluted soil in site 3 was able to grow on the medium containing the highest As^{5+} concentration
 289 (Table 3). It was selected for identification and further experiments.

290 The colony and micro-morphological features of the isolated fungal strain WKC-1 which was
 291 highly resistant to arsenic are presented in Figure S2 This strain was slow growing, achieving
 292 diameters of 22 to 45 mm in 21 days at 25 °C. The colonies appeared compact and dark greenish
 293 in colour. Under the microscope, the mycelium composed of septate, scarcely branched with
 294 thick-walled hyphae.

295
 296 The ITS rDNA sequence of the fungal isolate WKC-1 found in soil 3, conforms to phylogenetic
 297 lineage identical to the species *Acidomyces acidophilus*, CBS 335.97 (ex-type AJ 244237.1,
 298 FJ430711), which has previously been isolated from various highly acidic environments
 299 (Selbmann *et al.*, 2008; Hujsová *et al.*, 2013). The isolated fungal strain is designated *A.*
 300 *acidophilus* strain WKC-1 and has been given a GenBank accession number, KT727926 and the
 301 strain is deposited in DSMZ, Germany (DSM 105253) (Figure 2).




Figure 2: Phylogenetic dendrogram (scale bar = 5 represented nucleotide substitutions per 100 nucleotides. Numbers given at the nodes represent bootstrap values of 1000 replications).

Prior to the analysis of the isolated WKC-1 strain by MALDI TOF/TOF MS, the mass spectra of the three *A. acidophilus* reference strains were generated and inserted to the in-house database to create an *A. acidophilus* database library, since there is currently no database available for this species. The identification of *A. acidophilus* WKC-1 against three *A. acidophilus* reference strains showed that the isolated WKC-1 strain belongs to the *A. acidophilus* species with highly probable species identification to CBS 335.97 strain followed by secure genus identification against CCF 4251 and CCF 3679 strains (Figure 3).




Figure 3: Representation of mass spectra of isolated fungal strain WKC-1 and three *A. acidophilus* reference strains using MALDI-TOF/TOF MS.

*2.300-3.000 indicates high probable species identification; 2.000-2.299 indicates secure genus identification.

Table 3: The minimum inhibitory concentration ($\text{cm} \pm \text{standard error}$) of As^{5+} by *A. acidophilus* WKC-1 and the effect of low and high phosphate concentration. Data shown are the mean of three replicates.

Concentration of As ⁵⁺	Colony diameter (cm)	
	Medium containing 1 mg L ⁻¹ of PO ₄ ³⁻	Medium containing 100 mg L ⁻¹ of PO ₄ ³⁻
0	4.7 ± 0.3	5.3 ± 0.3***
1000	4.5 ± 0.2	4.9 ± 0.2**
7500	4.1 ± 0.2	4.5 ± 0.2*
15000	3.7 ± 0.2	4.3 ± 0.1***
20000	2.7 ± 0.1	4.0 ± 0.3***
22500	2.2 ± 0.2	3.6 ± 0.2***

324 Asterisks indicate statistical significance of differences tested by 2-sample t-test where * $p < 0.05$, ** $p <$
325 0.01, *** $p < 0.001$ compared to *A. acidophilus* WKC-1 containing 1 mg L⁻¹ of PO₄³⁻.

326 Minimum inhibitory concentration (MIC) of As for *A. acidophilus*

327 The MIC at pH 1 of isolated *A. acidophilus* WKC-1 reflects an extremely high tolerance for
328 arsenate, the strain could tolerate up to 22500 mg kg⁻¹ of As⁵⁺ in solid media. Two reference
329 strains of *A. acidophilus* (CBS 335.97 and CCF 4251) were tested for their tolerance to As⁵⁺ and
330 found to tolerate up to 10000 and 7500 mg L⁻¹ of As⁵⁺, respectively 2.5 times lower than the MIC
331 of the isolated WKC-1 strain (Table S3).


332

333 The CDA media containing 100 mg L⁻¹ of phosphate did have an effect on As⁵⁺ growth profile,
334 which resulted in increased resistance to As⁵⁺ (Table 3). The MIC between *A. acidophilus* WKC-
335 1 grown with 1 mg L⁻¹ of PO₄³⁻ and 100 mg L⁻¹ of PO₄³⁻ showed a statistical significant difference
336 in all media containing As⁵⁺ concentrations ranging from 1000 mg L⁻¹ to 22500 mg L⁻¹ ($p < 0.05$).

337

338 Effect of pH on fungal growth

339 Figure 4 presents the effect of pH on the growth characteristics of *A. acidophilus* WKC-1
340 colonies. The diameter of colony growth appeared to decrease as the pH increased and the *A.*
341 *acidophilus* WKC-1 strain can grow in extremely low pH of 1.



342

343 Figure 4: Colony diameter, representing a measurement of growth (cm \pm standard error) of
 344 isolated *A. acidophilus* WKC-1 at the minimum inhibitory concentration of As^{5+} , at different pH,
 345 at room temperature, on MEA. Data shown are the mean of three replicates.

346 * NG indicates no growth

347

348 **Arsenate removal efficiency**

349 In Figure 5, the mean percentages of arsenic removal by *A. acidophilus* WKC-1 and three *A.*
 350 *acidophilus* reference strains show that WKC-1 achieves a significantly higher percentage As^{5+}
 351 removal after 7, 14 and 21 days periods of cultivation compared to the *A. acidophilus* CBS
 352 335.97, CCF4251 and CCF3679 reference strains.




Figure 5: Percentage of arsenate removal by *A. acidophilus* WKC-1 and *A. acidophilus* reference strains after 7, 14 and 21 days cultivations of initial arsenate concentration of 100 mg L⁻¹. The error bars indicate the standard error of the mean of three replicates. Asterisks indicate statistical significance of differences tested by ANOVA where ** $p < 0.01$, *** $p < 0.001$ compared to *A. acidophilus* WKC-1.

There is a significant difference in As removal between the cultivation days ($p < 0.001$) for all four strains except for *A. acidophilus* CCF 4251, where there is no significant difference between 14 days and 21 days cultivation. The percentage removal of As⁵⁺ by *A. acidophilus* WKC-1 is 70.30 % after 21 days of cultivation compared to 56.30 %, 26.20 % and 25.80 % achieved with the reference strains CBS 335.97 and CCF 4251 and CCF 3679 respectively.

Biosorption of As

The summary of the effect of initial pH in the As⁵⁺ solution on the biosorption process of As⁵⁺ by *A. acidophilus* WKC-1 showed that there was an increase from 0.01 to 0.09 mg mg⁻¹ of the amount of As⁵⁺ absorbed by isolated *A. acidophilus* WKC-1 as the pH increased from 1.0 to 4.0 (Figure 6a). However, as the uptake started to decrease above pH 4.0, the optimum pH for the biosorption analysis of As⁵⁺ was set at pH 4.0.

373 The biomass loading with increased contact time was studied and it was found that the absorption
374 of As^{5+} rapidly increased in the first 30 min (Figure 6b). After 120 min, the sorption of As^{5+} by
375 *A. acidophilus* WKC-1 reached equilibrium and remained constant ($p > 0.05$). Therefore, the time
376 for the biosorption analysis for both As^{5+} and Sb^{5+} loaded biomass was set at 120 min. The effect
377 of biomass loading is summarized in Figure 6b. The sorption capacity by *A. acidophilus*
378 decreased as the biomass loading increased from 1 g L^{-1} to 5.0 g L^{-1} . In the presence of competing
379 Sb^{5+} ions, the As^{5+} uptake by fungal biomass is significantly affected ($p < 0.05$) as shown in Figure
380 6c.

381

382 The relationship between metalloid uptake capacity q_e , and equilibrium metal ion concentration
383 C_e , was evaluated based on the Langmuir model. The data from current study fitted the Langmuir
384 isotherm model well, with regression coefficient (R^2) of 0.989 (Figure 6d). Small b values (0.01)
385 imply strong binding of arsenic ions to *A. acidophilus* WKC-1. The predicted maximum capacity
386 of fungal strain uptake of As^{5+} by *A. acidophilus* WKC-1 was 170.82 mg g^{-1} dry biomass.

387




Figure 6. Biosorption of As^{5+} by *A. acidophilus* WKC-1; (a) the effects of pH on biomass loading; (b) the effect of contact time on As^{5+} biosorption at different concentration; (c) the effect of As^{5+} uptake in the presence and absence of Sb^{5+} ; and (d) the Langmuir isotherm plot of As^{5+} biosorption by *A. acidophilus* WKC-1 in the presence and absence of As^{5+} .

FT-IR analysis

FT-IR spectrum range of 4000-400 cm^{-1} was used to detect vibration frequency of changes in the functional group of isolated *A. acidophilus* strain before and after As^{5+} and Sb^{5+} loading (Figure 7). For control biomass spectrum (vibrational frequencies of bio-molecular functional groups), a broad band at 3306.55 cm^{-1} indicates -OH bonds stretching vibration at high concentration and

399 weak to medium of the -NH stretching (secondary amines). The peaks appearing in the 2921.74
400 and 2852.67 cm^{-1} region can be attributed to the strong asymmetric and symmetric stretching
401 vibration of CH_2 , respectively. Strong stretching vibrations of $\text{C}=\text{O}$ (esters) and $\text{C}=\text{O}$ (amide I
402 band) observed at peak 1744.18 and 1640.06 cm^{-1} respectively. The peak at 1640.06 cm^{-1} also
403 indicated variable symmetric stretching variations of $\text{C}=\text{C}$. The peak at 1544.68 cm^{-1} was
404 assigned to a motion of -NH bending (amide II) while the peak at 1456.46 cm^{-1} indicated medium
405 CH_2 and CH_3 deformation. O-H bending (in-plane) and strong stretching vibrations of C-F
406 appeared at the peak 1373.94 cm^{-1} . Medium to strong stretching vibrations of C-O and medium
407 C-N stretching of amine groups was observed at both 1238.95 and 1148.40 cm^{-1} peak.

408

409 A strong peak at 1028.03 cm^{-1} indicates P-OR (esters) as well as Si-OR groups. A NH_2 and N-H
410 wagging (shifts on H-bonding), C-H bending and ring puckering and a strong $=\text{C}-\text{H}$ & $=\text{CH}_2$
411 bending was observed at peak 887.86 cm^{-1} . The 'finger print' zone of the spectra, ranging from
412 500-700 cm^{-1} usually represents phosphate or sulfur functional groups.

413

414 For both As^{5+} and Sb^{5+} loaded spectra, significant shifts (weak/strong) were observed at
415 absorbance peaks 3306.55 cm^{-1} , 3003.79 cm^{-1} , 1373.94 cm^{-1} , 1028.03 cm^{-1} and 612.58 cm^{-1} either
416 by stretching vibrations, formation of new absorbance peaks and sharpening or lowering of the
417 shoulder peaks. These are the functional groups of -OH, -NH, -CH, - SO_3 , P-OR(esters) and PO_4 .

418




Figure 7: FT-IR spectra of *A. acidophilus* WKC-1 biomass (a) control, (b) As⁵⁺ loading and (c) Sb⁵⁺ loading. **Bold** indicates strong spectra shifting against the control.

423 Discussion

424 Soil abiotic characteristics and their interaction with the fungal isolates

425 Due to the igneous geology of Geevor tin mine, its activities generated various metal by-products
 426 such as Zn, Cu, As and Sb (Adriano, 1986; Hamilton, 2000). A process called roasting using a
 427 Brunton Calciner (burning furnace) where cassiterite (tin ore) containing As, Sb and other
 428 minerals such as Fe and Cu were burnt was used in Geevor tin mine. Large amount of roasting
 429 waste was deposited near to the production facilities. The contamination of As (Langdon *et al.*,
 430 2009) and Sb (Flynn *et al.*, 2003) found in the soil samples were most likely to come from the
 431 by-products of the roasting process used to obtain tin.

432

433 The three-step sequential extraction method provided information about the metals and
 434 metalloids potential mobility, bioavailability and amount bound to different soil fractions
 435 (Carapeto and Purchase, 2000; Lei *et al.*, 2010). This detail information is important for the
 436 evaluation of toxicity and bioavailability of metals and metalloids in soil from the mining dump
 437 as well as the feasibility of their remediation (Chen *et al.*, 2007). Total concentration of As and
 438 Sb were comparable to previously published data for mining sites in Cornwall by Peterson *et al.*
 439 (1979) and Dybowska *et al.* (2005) which presented concentration of As at 20 and 40 mm depth
 440 in the soils as high as 20,000 and 40000 mg As kg⁻¹, respectively. In addition, over 100 mg kg⁻¹
 441 of Sb levels in the soil have been recorded in close proximity to where the mining operations
 442 were carried out in Derbyshire, England (Li and Thornton, 1993). Most of the As was found in
 443 the residual fraction, which is not readily available and the metalloids present in this fraction can
 444 be used as a measurement of the degree of environmental pollution in soil. The higher the metals
 445 present in this fraction, the lower the degree of pollution (Howari and Banat, 2001).

446

447 The sum of concentrations in exchangeable and weakly organically bound fractions can be used
 448 to determine the bioavailability of transitional metals and metalloids in soils (Carapeto and
 449 Purchase, 2000). Geevor tin mine soils also contained high level of iron between 30000 and
 450 270000 mg kg⁻¹ (results not shown). According to Drahota and Filippi (2009), acidic conditions
 451 (pH<6) with relative abundance of iron oxide (Fe-oxide) may decrease the bioavailability of As
 452 in the soil with the formation of iron arsenates such as scorodites and pharmacosiderite in the
 453 mining soils (Jacobs *et al.*, 1970).

454

455 Identification of the isolated *A. acidophilus* WKC-1

456 *Acidomyces acidophilus* (Selbmann *et al.*, 2008) was first isolated by Starkey and Waksman
 457 (1943) in an extremely acidic, sulphate containing industrial water. Subsequently, more
 458 *Acidomyces acidomyces* strains were isolated in various extreme environments such as acid
 459 drainage (Germany), soil near sulfur pile (Canada), volcanic soil (Iceland), acidic industrial water
 460 (The Netherlands) and acid mine drainage water (USA) (Selbmann *et al.*, 2008). The
 461 morphologies of *Acidomyces* species were not easily described using microscopy because of its
 462 tendency to convert to meristematic growth, produce reluctantly disarticulating clumps of cells,
 463 or tend to appear to be entirely hyphal without any conidiation (Selbmann *et al.*, 2005; Selbmann
 464 *et al.*, 2008; Hujslová *et al.*, 2013).

465

466 *A. acidophilus* WKC-1 was identified by DNA sequencing and by MALDI-TOF/TOF MS. The
 467 latter is a robust method that is widely used in the identification of fungal species, especially
 468 clinical strains (Nenoff *et al.*, 2013). The growth period of *A. acidophilus* WKC-1 was
 469 significantly reduced (from 28 to 3 days) by culturing the fungal strain in liquid medium and
 470 incubating on a tube rotator. In order to obtain a trustworthy positive identification, the culture
 471 period for fungi should be no more than 10 days (De Respinis *et al.*, 2013). Since *A. acidophilus*
 472 is a black fungus, the pigment from the strain could inhibit the analysis using MALDI-TOF/TOF
 473 MS as the pigment will generate noise to the spectra produced (Buskirk *et al.*, 2011). However,
 474 such inhibition of obtaining spectra was not observed during the identification analysis.

475

476 *Penicillium* species was successfully identified using MALDI-TOF MS by Chen and Chen (2005)
 477 directly from intact fungal spores. Hettick *et al.*, (2008) obtained abundant peaks in the range
 478 5000-20000 m/z by using bead beating in the extraction process, the fungal samples and the
 479 MALDI-TOF MS in their experiment have identified all the 12 *Penicillium* species correctly.
 480 This study also show that the MALDI-TOF/TOF MS is a robust, cost saving and powerful system
 481 in fungal identification and characterization as suggested by Wieser *et al.* (2012).

482

483 However, the use of MALDI-TOF for identification of fungi has a few limitations. The spectral
 484 signal generated by MALDI-TOF is strongly influenced by the fungal growth medium as well as
 485 the protein extraction methods (Santos *et al.*, 2010). Due to the cell wall structure of fungi, protein
 486 extraction requires an additional step such as bead beating, to yield high quality spectra that

enable a valid identification (Croxatto *et al.*, 2012). The lack of reference spectra available in the database is the main disadvantage in using MALDI-TOF MS to identify fungal species and work like this current study can contribute to the development of a fungal database. The use of MALDI-TOF/TOF MS described in this paper has demonstrated that this method is capable to identify environmental fungi species provided that the correct sample preparation methods are being used.

Tolerance and removal efficiency of As

The soil condition where *A. acidophilus* WKC-1 was isolated is extremely hostile and inhabitable to most living organisms. However, the extreme acidity (pH 1) in the soil is a crucial factor for the growth of this acidophile. The ability of *A. acidophilus* to resist and survive in such acid and toxic environment is thought to be due to the presence and protection of a melanin-containing cell wall (Martin *et al.*, 1990).

A. acidophilus WKC-1 exhibits high As⁵⁺ removal efficiency even in extreme pH conditions. This indicates that *A. acidophilus* WKC-1 has great cellular detoxification mechanisms in toxic metalloids tolerance. The unique composition of fungal cell wall containing excellent metal-binding properties offers great advantage in metal removal either by entrapment in extra-cellular capsules and precipitation of metals (Gupta *et al.*, 2000). Previous study by Su *et al.* (2010) observed the intracellular uptake of As⁵⁺ in *Tichoderma asperellum* and *Fusarium oxysporum* can be as efficient as extracellular sorption in many fungi where the intracellular As⁵⁺ accumulation accounted for 82.2% and 63.4% of the total accumulated As⁵⁺.

Biosorption of As

The As⁵⁺ uptake by fungal biomass is significantly affected by the presence of competing ions, in this case Sb⁵⁺ (Figure 6). These ions compete for active binding sites due to the non-specificity of the functional groups present on the fungal cell surface. As a result, it is often found that specific transitional metal/metalloid(s) uptake from mixed solutions is lower than those in a solution containing the single transitional metal/metalloid.

The pH has profound effect on As⁵⁺ uptake by *A. acidophilus* WKC-1. The *A. acidophilus* WKC-1 As⁵⁺ sorption capacity increased with increasing pH from 1 to 4 and showed optimum As⁵⁺

adsorptions at pH 4 (Figure 6a). The pH of the solution affects the solubility of metalloid ions and the ionization state of the functional groups on the fungal cell wall by either interfering or enhancing with biosorption process (Fourest and Roux, 1992; Lopez *et al.*, 2000; Bayramoğlu *et al.*, 2003). Absorption of As^{5+} by WKC-1 at low pH is noticeably lower than at higher pH, this might be due to the competition between As^{5+} and H^+ or H_3O^+ ions present in the solution, for the negatively charged biosorbent binding sites (Gadd, 1994). It is likely that the high mobility and concentration of H^+ ions are preferentially adsorbed by the fungi cells than the studied metalloid ions. As the pH increases and the H^+ ion concentration in the solution decreases, a greater number of ligands (such as carboxyl, sulphhydryl, phosphate groups) with negative charges become available, thus increasing biosorption capacity (Feng *et al.*, 2011).

Higher absorption of As^{5+} was observed with increased contact time due to the abundant binding sites available on the fungal cell surface for the metal sorption by *A. acidophilus* WKC-1. The biomass loading of *A. acidophilus* WKC-1 for As^{5+} sorption was found to be optimal at 1 mg L^{-1} . The optimum biomass loading results support the hypothesis by Gupta and Rastogi (2008) that an increase in biomass loading could exert a shell effect by protecting the active binding sites from being occupied by the metal, resulting in the decrease of metal sorption. A similar effect of high biomass loading resulting in low sorption was observed by Bishnoi *et al.* (2007) in Cr (VI) removal by *Trichoderma viride*. In the presence of competing ions, metal uptake from mixed solutions is often found to be lower than those in a single-species system (Chong and Volesky, 1995).

In general, metal uptake by fungus increases as the ionic radius of the metal cation increases, thus metals with higher ionic charge show greater binding to biomass. However, as the concentration of other competing metalloid cations present within the same biosorption process increases, the uptake of another metalloid further decreases. Chemical interactions between two metal species as well as biomass may take place, resulting in competition for sorption sites on the surface (Akar *et al.*, 2005). Sari and Tuzen (2009) reported that maximum biosorption capacity of As^{5+} by *Inonotus hispidus* biomass was found to be 59.6 mg g^{-1} . Plant biomass prepared from sawdust of *Picea abies* has the maximum As^{5+} sorption capacity of 1.369 mg g^{-1} (Urik *et al.*, 2009). The As^{5+} adsorption capacity of zirconium (IV) loaded phosphoric chelate adsorbent, synthesized by radiation induced graft polymerization, was 149.8 mg g^{-1} (Seko *et al.*, 2004). *A. acidophilus* WKC-1 showed a greater As^{5+} adsorption capacity than previously studies fungal strains, where

the predicted maximum capacity of fungal strain uptake of As^{5+} by WKC-1 was 170.82 mg g^{-1} dry biomass and has potential to be used in bioremediation transitional metals in soils. The fate of As^{5+} after being adsorbed into the fungal cell might be broken down to less toxic species by powerful secondary enzymes produce intracellularly by the fungal itself or undergo a number detoxification pathways within the fungal cell which include the reduction to As^{3+} by arsenate reductases, followed by exclusion or sequestration of As^{3+} (Sharples *et al.*, 2000; González-Chávez *et al.*, 2002). Further study is required to elucidate and understand the detoxification mechanisms of As^{5+} of *A. acidophilus*.

The effect of other group 15 elements and phosphate on As removal

The effect of Sb^{5+} in reducing the As^{5+} sorption could be due to the competition of active binding sites as shown in the FT-IR analysis. Benjamin and Leckie (1981) showed that the adsorption of cadmium, copper, zinc and lead on amorphous iron oxyhydroxide were reduced in the presence of all the metals at the same time, as the availability of the active sorption binding sites decreased, which also lead to a decrease in the apparent adsorption equilibrium constants.

Similarly in the As^{5+} resistance experiment it was observed that PO_4^{3-} reduces As^{5+} toxicity on *A. acidophilus* WKC-1. According to Hughes (2002), PO_4^{3-} and As^{5+} are both tetrahedral oxy-anions and have similarity between structure, synthesis and hydrolysis thus PO_4^{3-} can chemically mimics and acts as a substitute to As^{5+} in biochemical processes by incorporated into the metabolic pathways of *A. acidophilus* unlike the As removal process.

These shifts in absorbance peaks of -OH/-NH as well as in phosphorus functions could indicate that alcohols/phenols, carboxylic acids and its derivatives, amine II and phosphate groups could be vital sites for the binding of As^{5+} ions. In the spectra of the As^{5+} loaded biomass, the shoulder peaks of 2852.74 , 1745.18 and 1023.16 cm^{-1} became sharper. Such observations could indicate that these related functional groups could be involved during the biosorption process. As seen in the ‘fingerprint’ region of the As^{5+} loaded biomass, multiple sharp peaks can be seen compared to the non-treated biomass. It was also noted that the absorbance of this region was much lower than the control sample. Phosphate and sulphur functional groups are indicating a possible interaction of the As^{5+} during biosorption process.

Based on the spectra from FT-IR generated, it suggests that As^{5+} and Sb^{5+} might compete for the binding sites of OH, -NH, -CH, -SO₃ and PO₄ functional groups on the surface of the isolated *A. acidophilus* WKC-1 strain. Previous study by Dixon (1997) showed that As^{5+} reacting in a similar way as phosphate in which it has the ability to form ester linkages with hydroxyl groups. A study carried out by Parascondola (1977) found out that Group 15 elements in the periodic table in both pentavalent and trivalent state can interact with sulphur (formation of As-S complexes), thus this supports the analysis from the FT-IR analysis that -SO₃ functional group could involve as a binding site of As^{5+} and Sb^{5+} . Some other functional groups such as C-O, C-N and CH₂ may also compete to a lesser extent by these two metalloids to bind on the surface of *A. acidophilus* WKC-1.

Conclusions

In conclusion, metal analysis showed that 26.40% of As^{5+} is bioavailable in the soil samples at a level below the MIC of *A. acidophilus* WKC-1, suggesting a good potential to apply this strain to remediate As polluted soil. The presence of phosphate decreases the toxicity of As^{5+} whereas Sb^{5+} significantly reduces the As removal ability of WCK-1. The -OH, -NH, -CH, -SO₃, and PO₄ functional groups have been identified as the key competitive binding sites between As^{5+} and Sb^{5+} . The isolate WKC-1 showed a high resistance and high percentage As^{5+} removal, one of the highest reported in *A. acidophilus* species. Our study also demonstrated that MALDI-TOF/TOF MS could provide a faster and cheaper way to identify environmental fungal strains. The tolerance of the isolated *A. acidophilus* WKC-1 strain to low pH and high As concentration together with its capacity to remove approximately 170 mg As^{5+} per gram dry biomass, made it an potential candidate to be used in bioremediation of As.

References

- Abrahams PW, Thornton I (1987) Distribution and extent of land contaminated by arsenic and associated metals in mining regions of southwest England. Institution of Mining and Metallurgy Transactions Section B Applied Earth Science 96:13–138.
- Adriano D (1986) Trace elements in the terrestrial environment. Springer, New York.

- 614 Akar T, Tunali S, Kiran I (2005) *Botrytis cinerea* as a new fungal biosorbent for removal of Pb(II)
 615 from aqueous solutions. *Biochem Eng J* 25:227-235.
 616
- 617 Antonucci I, Gallo G, Limauro D, Contursi P, Ribeiro AL, Blesa A, Berenguer J, Bartolucci S,
 618 Fiorentino G. (2017) An ArsR/SmtB family member regulates arsenic resistance genes unusually
 619 arranged in *Thermus thermophilus* HB27. *Microb Biotechnol.* 10:1690-1701.
 620 <https://doi.org/10.1111/1751-7915.12761>.
 621
- 622 Asklund R, Eldvall B (2005) Contamination of water resources in Tarkwa mining area of Ghana.
 623 *Resource* 27:61-75.
 624
- 625 ASTM (2000) Standard test methods for moisture, ash, and organic matter of peat and other
 626 organic soils. Method D 2974-14. American Society for Testing and Materials. West
 627 Conshohocken, PA. <http://www.astm.org/Standards/D2974.htm>.
 628
- 629 Bayramoğlu G, Bektaş S, Arica MY (2003) Biosorption of heavy metal ions on immobilized
 630 white-rot fungus *Trametes versicolor*. *J Hazard Mater* 101:285-300.
 631
- 632 Benjamin MM, Leckie JO (1981) Multiple-site adsorption of Cd, Cu, Zn, and Pb on amorphous
 633 iron oxyhydroxide. *J Colloid Interface Sci* 79:209-221.
 634
- 635 Bishnoi NR, Kumar R, Bishnoi K (2007) Biosorption of Cr (VI) with *Trichoderma viride*
 636 immobilized fungal biomass and cell free Ca-alginate beads. *Indian J Exp Biol* 45:657.
 637
- 638 Buskirk AD, Hettick JM, Chipinda I, Law BF, Siegel PD, Slaven JE, Green BJ, Beezhold DH
 639 (2011) Fungal pigments inhibit the matrix-assisted laser desorption/ionization time-of-flight mass
 640 spectrometry analysis of darkly pigmented fungi. *Anal Biochem* 411:122-128.
 641
- 642 Camm GS, Powell N, Glass H, Cressey G, Kirk C (2003) Soil geochemical signature of a calciner
 643 site, Cornwall, SW England. *Appl Earth Sci* 112:268-278.
 644
- 645 Camm GS, Glass HJ, Bryce DW, Butcher AR (2004) Characterisation of a mining-related
 646 arsenic-contaminated site, Cornwall, UK. *J Geochem Explor* 82:1–15.
 647

- 648 Carapeto C, Purchase D (2000) Use of sequential extraction procedures for the analysis of
649 cadmium and lead in sediment samples from a constructed wetland. *Bull Environ Cont Toxicol*
650 64:51-58.
651
- 652 Chen HY, Chen YC (2005) Characterization of intact *Penicillium* spores by matrix-assisted laser
653 desorption/ionization mass spectrometry. *Rapid Comm Mass Spectrom* 19:3564-3568.
654
- 655 Chen Z, He M, Sakurai K, Kang Y, Iwasaki K (2007) Concentrations and chemical forms of
656 heavy metals in urban soils of Shanghai, China. *Soil Scie Plant Nutr* 53:517-529.
657
- 658 Chong K, Volesky B (1995) Description of two-metal biosorption equilibria by Langmuir-type
659 models. *Biotech Bioeng* 47:451-460.
660
- 661 Croxatto A, Prod'hom G, Greub G (2012) Applications of MALDI-TOF mass spectrometry in
662 clinical diagnostic microbiology. *FEMS Microbiol Rev* 36:380-407.
663
- 664 De Respinis S, Tonolla M, Pranghofer S, Petrini L, Petrini O, Bosshard PP (2013) Identification
665 of dermatophytes by matrix-assisted laser desorption/ionization time-of-flight mass
666 spectrometry. *Med Mycol* 51:514-521.
667
- 668 Defra (2014) Development of Category 4 Screening Levels for Assessment of Land Affected by
669 Contamination – SP1010. Department for Environment, Food and Rural Affairs.
670 <http://randd.defra.gov.uk/Default.aspx?Menu=Menu&Module=More&Location=None&Completed=0&ProjectID=18341>. Accessed 15 Nov 2016
671
672
- 673 Del Giudice I, Limauro D, Pedone E, Bartolucci S, Fiorentino G. (2013) A novel arsenate
674 reductase from the bacterium *Thermus thermophilus* HB27: its role in arsenic detoxification.
675 *Biochim Biophys Acta*. 1834:2071-2079. doi [10.1016/j.bbapap.2013.06.007](https://doi.org/10.1016/j.bbapap.2013.06.007).
676
- 677 Dixon HB (1997) The biochemical action of arsonic acids especially as phosphate analogues.
678 *Adv Inorg Chem* 44:191-227.
679
- 680 Dos Santos JV, de Melo Rangel W, Guimaraes AA, Jaramillo PMD, Rufini M, Marra LM, López
681 MV, Da Silva MAP, Soares CRFS, de Souza Moreira FM (2013) Soil biological attributes in

- 682 arsenic-contaminated gold mining sites after revegetation. *Ecotox* 22:1526-1537.
683
- 684 Drahotka P, Filippi M (2009) Secondary arsenic minerals in the environment: a review. *Environ*
685 *Int* 35:1243-1255.
686
- 687 Dutch Environment Ministry (2013) Annexes circular on target values and intervention values
688 for soil remediation. Dutch National Institute of Public Health & the Environment (RIVM), The
689 Netherlands
690
- 691 Dybowska A, Farago M, Valsami-Jones E, Thornton I (2005) Operationally defined associations
692 of arsenic and copper from soil and mine waste in south-west England. *Chem Speciation Bioavail*
693 17:147-160.
694
- 695 Feng N, Guo X, Liang S, Zhu Y, Liu J (2011) Biosorption of heavy metals from aqueous solutions
696 by chemically modified orange peel. *J Hazard Mater* 185:49-54.
697
- 698 Flynn HC, Meharg AA, Bowyer PK, Paton GI (2003) Antimony bioavailability in mine soils.
699 *Environ Poll* 124:93-100.
700
- 701 Fourest E, Roux JC (1992) Heavy metal biosorption by fungal mycelial by-products: mechanisms
702 and influence of pH. *Appl Microbiol Biotech* 37:399-403.
703
- 704 Francesconi K, Visoottiviseth P, Sridokchan W, Goessler W (2002) Arsenic species in an arsenic
705 hyperaccumulating fern, *Pityrogramma calomelanos*: a potential phytoremediator of arsenic-
706 contaminated soils. *Sci Total Environ* 284:27-35.
707
- 708 Gadd GM (2009) Biosorption: critical review of scientific rationale, environmental importance
709 and significance for pollution treatment. *J Chem Tech Biotech* 84:13-28.
710
- 711 Gadd GM (1994) Interactions of Fungi with Toxic Metals. In: Powell K.A., Renwick A., Peberdy
712 J.F. (eds) *The Genus Aspergillus*. Federation of European Microbiological Societies Symposium
713 Series, vol 69. Springer, Boston, MA.
714

- 715 Gillman G, Sumpter E (1986) Modification to the compulsive exchange method for measuring
716 exchange characteristics of soils. *Soil Res* 24:61-66.
717
- 718 Gonzalez-Chavez C, Harris PJ, Dodd J, Meharg AA (2002) Arbuscular mycorrhizal fungi confer
719 enhanced arsenate resistance on *Holcus lanatus*. *New Phytologist* 155:163-71.
720
- 721 Guibaud G, Tixier N, Bouju A, Baudu M (2003) Relation between extracellular polymers'
722 composition and its ability to complex Cd, Cu and Pb. *Chemosphere* 52:1701-1710.
723
- 724 Gupta R, Ahuja P, Khan S, Saxena RK, Mohapatra H (2000) Microbial biosorbents: Meeting
725 challenges of heavy metal pollution in aqueous solutions. *Curr Sci* 78:967-73.
726
- 727 Gupta VK, Rastogi A (2008) Biosorption of lead (II) from aqueous solutions by non-living algal
728 biomass *Oedogonium* sp. and *Nostoc* sp.—a comparative study. *Colloids Surf B* 64:170-178.
729
- 730 Hamilton E (2000) Environmental variables in a holistic evaluation of land contaminated by
731 historic mine wastes: a study of multi-element mine wastes in West Devon, England using arsenic
732 as an element of potential concern to human health. *Sci Total Environ* 249:171-221.
733
- 734 Hess M (2008) Thermoacidophilic proteins for biofuel production. *Trends Microbiol* 16:414-419.
735
- 736 Hettick JM, Green BJ, Buskirk AD, Kashon ML, Slaven JE, Janotka E, Blachere FM, Schmechel
737 D, Beezhold DH (2008) Discrimination of *Penicillium* isolates by matrix-assisted laser
738 desorption/ionization time-of-flight mass spectrometry fingerprinting. *Rapid Comm Mass*
739 *Spectrom* 22:2555-2560.
740
- 741 Howari F, Banat K (2001) Assessment of Fe, Zn, Cd, Hg, and Pb in the Jordan and Yarmouk
742 river sediments in relation to their physicochemical properties and sequential extraction
743 characterization. *Water Air Soil Pollut* 132:43-59.
744
- 745 Hudson-Edwards KA, Macklin M, Brewer P, Dennis I (2008) Environment Agency Science
746 Report SC030136/SR4 Assessment of metal mining-contaminated river sediments in England
747 and Wales. Environment Agency Bristol.

- 748
- 749 Hughes MF (2002) Arsenic toxicity and potential mechanisms of action. *Toxicol Lett* 133:1-6.
- 750
- 751 Hujslová M, Kubátová A, Kostovčík M, Kolařík M (2013) *Acidiella bohémica* gen. et sp. nov.
- 752 and *Acidomyces* spp.(Teratosphaeriaceae), the indigenous inhabitants of extremely acidic soils in
- 753 Europe. *Fungal Divers* 58:33-45.
- 754
- 755 Jacobs L, Syers J, Keeney D (1970) Arsenic sorption by soils. *Soil Sci Soc America J* 34:750-
- 756 754.
- 757
- 758 Jacobson ES, Hove E, Emery HS (1995) Antioxidant function of melanin in black fungi. *Infect*
- 759 *Immun* 63:4944-4945.
- 760
- 761 Jung WH, Sham A, White R, Kronstad JW (2006) Iron regulation of the major virulence factors
- 762 in the AIDS-associated pathogen *Cryptococcus neoformans*. *PLoS Biol* 4:e410.
- 763
- 764 Langdon C, Morgan A, Charnock J, Semple KT, Lowe C (2009) As-resistance in laboratory-
- 765 reared F1, F2 and F3 generation offspring of the earthworm *Lumbricus rubellus* inhabiting an
- 766 As-contaminated mine soil. *Environ Poll* 157:3114-3119.
- 767
- 768 Lei M, Zhang Y, Khan S, Qin PF, Liao BH (2010) Pollution, fractionation, and mobility of Pb,
- 769 Cd, Cu, and Zn in garden and paddy soils from a Pb/Zn mining area. *Environ Monitor Assess*
- 770 168:215-222.
- 771
- 772 Li X, Thornton I (1993) Arsenic, antimony and bismuth in soil and pasture herbage in some old
- 773 metalliferous mining areas in England. *Environ Geochem Health* 15:135-144.
- 774
- 775 Lopez A, Lazaro N, Priego J, Marques A (2000) Effect of pH on the biosorption of nickel and
- 776 other heavy metals by *Pseudomonas fluorescens* 4F39. *J Industrial Microbiol Biotech* 24:146-
- 777 151.
- 778
- 779 Martin AM, Chintalapati SP, Patel TR (1990) Extraction of bitumens and humic substances from
- 780 peat and their effects on the growth of an acid-tolerant fungus. *Soil Biol Biochem* 22:949-954.
- 781

- 782 Martin KJ, Rygiewicz PT (2005) Fungal-specific PCR primers developed for analysis of the ITS
783 region of environmental DNA extracts. *BMC Microbiol* 5:1.
784
- 785 Nenoff P, Erhard M, Simon JC, Muylowa GK, Herrmann J, Rataj W, Gräser Y (2013) MALDI-
786 TOF mass spectrometry—a rapid method for the identification of dermatophyte species. *Med*
787 *Mycol* 51:17-24.
788
- 789 Noall C (1973) The St Just Mining District. Bradford Barton, Truro, pp. 179.
790
- 791 Olaniran AO, Balgobind A, Pillay B (2013) Bioavailability of heavy metals in soil: impact on
792 microbial biodegradation of organic compounds and possible improvement strategies. *Int J Mol*
793 *Sci* 15:10197-228.
794
- 795 Peijnenburg W, Jager T (2003) Monitoring approaches to assess bioaccessibility and
796 bioavailability of metals: matrix issues. *Ecotox Environ Safety* 56:63-77.
797
- 798 Peterson PJ, Benson LM, Porter EK (1979) Biogeochemistry of arsenic on polluted sites in SW
799 England. In: International conference of Management and Control of Heavy Metals in the
800 Environment. CEP Consultants Ltd, Edinburgh, pp 198–201.
801
- 802 Pirrie D, Power MR, Wheeler PD, Cundy A, Bridges C, Davey G (2002) Geochemical signature
803 of historical mining: Fowey Estuary, Cornwall, UK. *J Geochem Explor* 76:31-43.
804
- 805 Polizeli M, Rizzatti A, Monti R, Terenzi H, Jorge JA, Amorim D (2005) Xylanases from fungi:
806 properties and industrial applications. *Appl Microbiol Biotech* 67:577-591.
807
- 808 Razo I, Carrizales L, Castro J, Díaz-Barriga F, Monroy M (2004) Arsenic and heavy metal
809 pollution of soil, water and sediments in a semi-arid climate mining area in Mexico. *Water Air*
810 *Soil Pollut* 152:129-152.
811
- 812 Santos C, Paterson RRM, Venâncio A, Lima N (2010) Filamentous fungal characterizations by
813 matrix assisted laser desorption/ionization time-of-flight mass spectrometry. *J Appl Microbiol*
814 108:375-385.

- 815
- 816 Sari A, Tuzen M (2009) Biosorption of As(III) and As(V) from aqueous solution by macrofungus
 817 (*Inonotus hispidus*) biomass: equilibrium and kinetic studies. J Hazard Mater 164:1372-1378.
 818
- 819 Seko N, Basuki F, Tamada M, Yoshii F (2004) Rapid removal of arsenic (V) by zirconium (IV)
 820 loaded phosphoric chelate adsorbent synthesized by radiation induced graft polymerization.
 821 React Funct Polym 59:235-241.
 822
- 823 Selbmann L, De Hoog G, Mazzaglia A, Friedmann E, Onofri S (2005) Fungi at the edge of life:
 824 cryptoendolithic black fungi from Antarctic desert. Stud Mycol 51:1-32.
 825
- 826 Selbmann L, De Hoog GS, Zucconi L, Isola D, Ruisi S, van den Ende AG, Ruibal C, De Leo F,
 827 Urzi C, Onofri S (2008) Drought meets acid: three new genera in a dothidealean clade of
 828 extremotolerant fungi. Stud Mycol 61:1-20.
 829
- 830 Sharples JM, Meharg AA, Chambers SM, Cairney JW (2000) Mechanism of arsenate resistance
 831 in the ericoid mycorrhizal fungus *Hymenoscyphus ericae*. Plant Physiol 124:1327-34.
 832
- 833 Sigler L, Carmichael J (1974) A new acidophilic *Scytalidium*. Can J Microbiol 20:267-268.
 834
- 835 Starkey RL, Waksman SA (1943) Fungi tolerant to extreme acidity and high concentrations of
 836 copper sulfate. J Bacteriol 45:509.
 837
- 838 Stirling D (2003) DNA extraction from fungi, yeast, and bacteria. PCR Protocols vol:53-54.
 839
- 840 Su S, Zeng X, Bai L, Jiang X, Li L (2010) Bioaccumulation and biovolatilisation of pentavalent
 841 arsenic by *Penicillium janthinellum*, *Fusarium oxysporum* and *Trichoderma asperellum* under
 842 laboratory conditions. Curr Microbiol 61:261-6.
 843
- 844 Tamura K, Stecher G, Peterson D, Filipski A, Kumar S (2013) MEGA6: molecular evolutionary
 845 genetics analysis version 6.0. Mol Biol Evol 30:2725-2729.
 846
- 847 Telford K, Maher W, Krikowa F, Foster S, Ellwood MJ, Ashley PM, Lockwood PV, Wilson SC

- 848 (2009) Bioaccumulation of antimony and arsenic in a highly contaminated stream adjacent to the
849 Hillgrove Mine, NSW, Australia. *Environ Chem* 6:133-143.
- 850
- 851 Tetsch L, Bend J, Hölker U (2006) Molecular and enzymatic characterisation of extra-and
852 intracellular laccases from the acidophilic ascomycete *Hortaea acidophila*. *Antonie Van*
853 *Leeuwenhoek* 90:183-194. Doi: <https://doi.org/10.1007/s10482-006-9064-z>
- 854
- 855 Tsezos M, Remoudaki E, Angelatou V (1996) A study of the effects of competing ions on the
856 biosorption of metals. *Int Biodeterior Biodegradation* 38:19-29.
- 857
- 858 Urik M, Littera P, Kolen M (2009) Removal of arsenic (V) from aqueous solutions using
859 chemically modified sawdust of spruce (*Picea abies*): kinetics and isotherm studies. *Int J Environ*
860 *Sci Technol* 6:451-456.
- 861
- 862 Van Veen E, Lottermoser B, Parbhakar-Fox A, Fox N, Hunt J (2016) A new test for plant
863 bioaccessibility in sulphidic wastes and soils: a case study from the Wheal Maid historic tailings
864 repository in Cornwall, UK. *Sci Total Environ* 563:835-844.
- 865
- 866 Vijayaraghavan K, Padmesh T, Palanivelu K, Velan M (2006) Biosorption of nickel (II) ions onto
867 *Sargassum wightii*: application of two-parameter and three-parameter isotherm models. *J Hazard*
868 *Mater* 133:304-308.
- 869
- 870 Wang S, Mulligan CN (2006) Occurrence of arsenic contamination in Canada: sources, behavior
871 and distribution. *Sci Total Environ* 366:701-721.
- 872
- 873 White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal
874 ribosomal RNA genes for phylogenetics. *PCR Protocol* 18:315-322.
- 875
- 876 Wieser A, Schneider L, Jung J, Schubert S (2012) MALDI-TOF MS in microbiological
877 diagnostics—identification of microorganisms and beyond (mini review). *Appl Microbiol*
878 *Biotech* 93:965-974.
- 879
- 880 Xu X, Xia L, Huang Q, Gu JD, Chen W (2012) Biosorption of cadmium by a metal-resistant
881 filamentous fungus isolated from chicken manure compost. *Environ Technol* 33:1661-1670.

- 882
- 883 Yim W-S (1981) Geochemical investigations on fluvial sediments contaminated by tin-mine
- 884 tailings, Cornwall, England. *Environ Geol* 3:245-256.




Fig. S1: An aerial photograph of the Geevor tin-mine in Pendene, Penzance, Cornwall, UK and locations of soil sampling sites (Greevor Tin Mine was viewed on 17 July 2013. <https://www.google.co.uk/maps/places/Greevor+Tin+Mine/@50.1519033,-5.6744307,1404m>).




Figure S2: Morphological features the fungus of (a) colony of the isolated fungal strain in CDA medium, (b) Hyphae of the strain observed by light microscope at a magnification of 400x and (c) scanning electron microscope (SEM) at a magnification of 1000x (b) and 2200x (c), scale bar in (b) and (c) = 2 μ m.

892 Table S1: Operating parameters of ICP- OES (iCAP 1600)

Operating parameters of the thermos ICP-OES (iCAP 1600)	
Power (W)	1150
Auxiliary gas flow (L/min)	0.5
Nebuliser gas flow (L/min)	0.75
Coolant gas flow(L/min)	12
View	Axial
Purge gas flow	Normal
Flush pump rate (rpm)	100
Analysis pump rate (rpm)	50
Camera temperature	-47
Optics temperature	38

893

894 Table S2: Recovery of As and Sb (mg kg^{-1}) metal using certified reference material, SRM 2710a
 895 Montana Soil using acid digestion method. Data shown are the mean of three replicates.

Element	Certified value	Mean obtained value	Average % recovery
Reference material Lot 011233			
As	61.10 ± 2.08	57.05 ± 0.20	93.37
Sb	73.7 ± 10.50	62.19 ± 2.03	84.38
Reference material Lot 017309			
As	202.00 ± 17.70	186.30 ± 12.32	92.23
Sb	125.00 ± 13.53	109.9 ± 12.90	87.92

896 Table S3: The diameter measurement of the minimum inhibitory concentration of As⁵⁺ by
 897 isolated *A. acidophilus* and two positive control *A. acidophilus* type strains

As ⁵⁺ concentration	Diameter (cm)		
	Isolated <i>A. acidophilus</i> strain	Positive control	
		<i>A. acidophilus</i>	<i>A. acidophilus</i>
		CBS 335.07	CBS 4251
Control	4.7 ± 0.2	3.9 ± 0.4	4.2 ± 0.2
1000	4.5 ± 0.1	3.8 ± 0.1	3.9 ± 0.4
7500	4.3 ± 0.2	2.7 ± 0.2	2.4 ± 0.1
10000	4.1 ± 0.2	1.4 ± 0.1	NG
12500	3.9 ± 0.1	NG	NG
15000	3.7 ± 0.3	NG	NG
17500	3.4 ± 0.1	NG	NG
20000	2.7 ± 0.2	NG	NG
22500	2.2 ± 0.2	NG	NG
25000	NG	NG	NG

898 * NG indicates no growth